

文章编号:1005-3085(2011)03-0335-08

一类2-指标变延迟微分代数方程BDF方法的收敛性*

刘红良, 肖爱国

(湘潭大学数学与计算科学学院, 科学与工程计算与数值仿真湖南省重点实验室, 湘潭 411105)

摘 要: 延迟微分代数方程经常出现在自动控制、电力和电路分析、多体动力学等许多实际问题中. 目前对延迟微分代数方程数值分析研究主要集中于线性问题和1-指标问题; 对高指标非线性延迟微分代数方程数值分析的研究较困难, 国内外仅有少量工作且大多为常延迟. 本文将向后微分公式(BDF)应用于求解2-指标非线性变延迟微分代数方程, 获得了相应的收敛性结果, 并通过数值试验进行了验证.

关键词: 2-指标微分代数方程; 向后微分公式; 收敛性; 变延迟

分类号: AMS(2000) 90C29

中图分类号: O221.4

文献标识码: A

1 引言

目前, 对延迟微分代数方程的研究思想许多借助了微分代数方程的理论及延迟微分方程的理论, 因此常微分方程, 微分代数方程及(中立型)延迟微分方程^[1,2]等常用的理论及数值方法被许多学者应用于延迟微分代数方程及其数值方法. 但是, 延迟微分代数方程因有延迟的影响, 很多特性是微分代数方程或延迟微分方程所没有的, 所以对延迟微分代数方程的研究必须独立进行. Petzold^[3]指出“微分代数方程不是常微分方程”, 同样, 延迟微分代数方程既不是微分代数方程, 也不是延迟微分方程.

迄今为止, 国内外对延迟微分代数方程的研究集中在理论分析及数值方法的稳定性和误差分析上. 在理论分析方面, 重点研究了(中立型)延迟(积分)微分代数方程的结构分析及渐近稳定性^[4-7]. 在数值稳定性方面, 重点讨论了Rosenbrock、(多步)Runge-Kutta(RK)方法及线性多步法(含 θ -方法、BDF方法)关于线性(中立型)常延迟(积分)微分代数方程的渐近稳定性^[4,5,8,9]及 θ -方法(含中点法)关于1-指标刚性延迟微分代数方程的稳定性^[10,11]. 在数值误差分析方面, 文献[12-14]分别获得了RK方法、线性多步方法、单支方法及块方法关于1-指标、2-指标常延迟微分代数方程的误差分析结果, 文献[15]获得了配置方法关于1-指标、2-指标变延迟半显式非线性微分代数系统的收敛性结果, 文献[16,17]获得了单支方法和RK方法关于1-指标变延迟微分代数方程的收敛性结果.

本文应用向后微分公式(BDF)求解一类2-指标非线性变延迟微分代数方程, 获得了该方法的收敛性结果, 并通过数值试验进行了验证.

收稿日期: 2009-08-05. 作者简介: 刘红良(1979年11月生), 男, 博士. 研究方向: 刚性微分方程数值方法.

*基金项目: 国家自然科学基金(10971175); 教育部高校博士点基金(20094301110001); 湖南省自然科学基金(09JJ3002); 湘潭大学博士科研启动项目(11QDZ01).

2 一类2-指标变延迟微分代数方程的收敛性分析

考虑一类2-指标变延迟微分代数方程(2-DDAEs)

$$\begin{cases} y'(x) = f(y(x), y(x - \tau(x)), z(x)), & x \in [0, T], \\ 0 = g(y(x)), & x \in [0, T], \\ z(0) = z_0, \quad y(x) = \varphi(x), & x \in [-\tau, 0], \end{cases} \quad (1)$$

这里延迟函数 $\tau(x)$ 可导, 且满足 $0 < \tau(x) \leq \tau$, $0 < \tau'(x) < 1$, 函数 $f: R^{n_1} \times R^{n_1} \times R^{n_2} \rightarrow R^{n_1}$, $g: R^{n_1} \rightarrow R^{n_2}$ 充分光滑且各阶导数有界, 初值函数 $\varphi(x)$ 连续, 且 $g_y(y)f_z(y, y(x - \tau(x)), z)$ 可逆. 恒设问题 (1) 有唯一光滑真解 $y(x), z(x)$. 在下文中符号 $\|\cdot\|$ 表示欧氏范数.

把带 p 阶插值公式的 BDF 方法应用于求解问题 (1), 可得

$$\sum_{i=0}^k \alpha_i y_{n+i} = hf(y_{n+k}, y_n^h, z_{n+k}), \quad 0 = g(y_{n+k}), \quad (2)$$

这里 $x_{n+k} = x_n + kh$, $n \geq 0$, 且

$$y_n^h = \begin{cases} \varphi(x_{n+k} - \tau(x_{n+k})), & x_{n+k} - \tau(x_{n+k}) \leq 0, \\ \sum_{j=-u}^q Q_j(\delta_n) y_{n+k-m_n+j}, & x_{n+k} - \tau(x_{n+k}) > 0, \end{cases} \quad (3)$$

其中 $\tau(x_{n+k}) = (m_n - \delta_n)h$, $u, q, m_n \in Z^+$, $\delta_n \in [0, 1]$, $q + u = p$, $q + 1 \leq m_n$, $Q_j(\delta_n)$ 为拉格朗日插值基函数.

考虑 (2) 的扰动问题

$$\sum_{i=0}^k \alpha_i \hat{y}_{n+i} = hf(\hat{y}_{n+k}, \hat{y}_n^h, \hat{z}_{n+k}) + h\delta, \quad 0 = g(\hat{y}_{n+k}) + \theta. \quad (4)$$

定理 1 设 y_{n+k}, z_{n+k} 由 (2) 确定, 而扰动 $\hat{y}_{n+k}, \hat{z}_{n+k}$ 由 (4) 确定, 起始值满足

$$\begin{aligned} y_{n+j} - y(x_{n+j}) &= O(h), \quad z_{n+j} - z(x_{n+j}) = O(h), \\ g(y_{n+j}) &= O(h^2), \quad j = 0, 1, \dots, k-1, \quad x_{n+j} = x_n + jh, \end{aligned} \quad (5)$$

及起始值扰动满足

$$\hat{y}_{n+j} - y_{n+j} = O(h^2), \quad \hat{z}_{n+j} - z_{n+j} = O(h), \quad \delta = O(h), \quad \theta = O(h^2), \quad (6)$$

则对 $h < h_0$, 有

$$\|\hat{y}_{n+k} - y_{n+k}\| \leq C(\|\hat{Y}_n - Y_n\| + h\|\delta\| + \|\theta\|), \quad (7)$$

$$\|\hat{z}_{n+k} - z_{n+k}\| \leq \frac{C}{h} \left(\sum_{j=0}^{k-1} \|g_y(\hat{y}_{n+k})(\hat{y}_{n+j} - y_{n+j})\| + h\|\hat{Y}_n - Y_n\| + h\|\delta\| + \|\theta\| \right), \quad (8)$$

这里

$$\hat{Y}_n - Y_n = (\hat{y}_{n+k-1}^T - y_{n+k-1}^T, \dots, \hat{y}_n^T - y_n^T, (\hat{y}_n^h)^T - (y_n^h)^T)^T, \quad (9)$$

$$\|\hat{Y}_n - Y_n\| = \max \left(\max_{0 \leq j \leq k-1} \|\hat{y}_{n+j}^T - y_{n+j}^T\|, \|(\hat{y}_n^h)^T - (y_n^h)^T\| \right). \quad (10)$$

证明 令

$$\eta = -\sum_{i=0}^{k-1} \frac{\alpha_i}{\alpha_k} y_{n+i}, \quad \hat{\eta} = -\sum_{i=0}^{k-1} \frac{\alpha_i}{\alpha_k} \hat{y}_{n+i}. \quad (11)$$

调整 h 和 δ 比例, 使得 (2) 和 (4) 分别变为

$$y_{n+k} = \eta + hf(y_{n+k}, y_n^h, z_{n+k}), \quad 0 = g(y_{n+k}), \quad (12)$$

$$\hat{y}_{n+k} = \hat{\eta} + hf(\hat{y}_{n+k}, \hat{y}_n^h, \hat{z}_{n+k}) + h\delta, \quad 0 = g(\hat{y}_{n+k}) + \theta. \quad (13)$$

(12) 中第二式可改写为

$$0 = g(y_{n+k}) - g(\eta) + g(\eta) = \int_0^1 g_y(\eta + \tau(y_{n+k} - \eta)) d\tau [y_{n+k} - \eta] + g(\eta), \quad (14)$$

把 (12) 中第一式代入 (14) 可得

$$\int_0^1 g_y(\eta + \tau(y_{n+k} - \eta)) d\tau f(y_{n+k}, y_n^h, z_{n+k}) + \frac{1}{h} g(\eta) = 0, \quad (15)$$

同理, (13) 中第二式改写为

$$\int_0^1 g_y(\hat{\eta} + \tau(\hat{y}_{n+k} - \hat{\eta})) d\tau [f(\hat{y}_{n+k}, \hat{y}_n^h, \hat{z}_{n+k}) + \delta] + \frac{1}{h} g(\hat{\eta}) + \frac{1}{h} \theta = 0. \quad (16)$$

利用函数 f, g 的光滑性及 $g_y f_z$ 的可逆性, 由 (15)-(16) 可得

$$\begin{aligned} \|\hat{z}_{n+k} - z_{n+k}\| \leq C_1 & \left(\|\hat{y}_{n+k} - y_{n+k}\| + \|\hat{\eta} - \eta\| + \|\hat{y}_n^h - y_n^h\| \right. \\ & \left. + \|\delta\| + \frac{1}{h} \|\theta\| + \frac{1}{h} \|g(\hat{\eta}) - g(\eta)\| \right), \end{aligned} \quad (17)$$

由 (12) 中第一式减去 (13) 中第一式可得

$$\begin{aligned} \|\hat{y}_{n+k} - y_{n+k}\| & \leq \|\hat{\eta} - \eta\| + h \|f(\hat{y}_{n+k}, \hat{y}_n^h, \hat{z}_{n+k}) - f(y_{n+k}, y_n^h, z_{n+k})\| \\ & \leq \|\hat{\eta} - \eta\| + hL (\|\hat{y}_{n+k} - y_{n+k}\| + \|\hat{y}_n^h - y_n^h\| + \|\hat{z}_{n+k} - z_{n+k}\|), \end{aligned} \quad (18)$$

这里 L 为右端函数 f 的经典 Lipschitz 常数, 把 (17) 代入 (18), 化简得

$$\|\hat{y}_{n+k} - y_{n+k}\| \leq C_2 (\|\hat{\eta} - \eta\| + \|\hat{y}_n^h - y_n^h\| + h\|\delta\| + \|\theta\|), \quad h \leq \frac{1}{L + LC_1}. \quad (19)$$

把 (19) 代入 (17), 化简得

$$\begin{aligned} \|\hat{z}_{n+k} - z_{n+k}\| & \leq C_3 \left(\|\hat{\eta} - \eta\| + \|\hat{y}_n^h - y_n^h\| + \|\delta\| \right. \\ & \left. + \frac{1}{h} \|\theta\| + \frac{1}{h} \|g_y(\hat{\eta})(\hat{\eta} - \eta)\| \right), \quad h \leq \frac{1}{L + LC_1}, \end{aligned} \quad (20)$$

又

$$\|\hat{\eta} - \eta\| = \left\| \sum_{j=0}^{k-1} \frac{\alpha_j}{\alpha_k} (\hat{y}_{n+j} - y_{n+j}) \right\| \leq C_4 \|\hat{Y}_n - Y_n\|, \quad (21)$$

$$\begin{aligned} \|g_y(\hat{\eta})(\hat{\eta} - \eta)\| & \leq \|g_y(\hat{y}_k)(\hat{\eta} - \eta)\| + \|\hat{\eta} - \eta\| O(h) \\ & \leq C_5 \left(\sum_{j=0}^{k-1} \|g_y(\hat{y}_k)(\hat{y}_{n+j} - y_{n+j})\| + \|\hat{Y}_n - Y_n\| O(h) \right). \end{aligned} \quad (22)$$

把(21), (22)代入(19), (20), 经化简得结论(7), (8).

推论 1 设2-DDAEs (1) 满足 $g_y f_z$ 可逆, 且带 p 阶插值公式 (3) 的 BDF 方法 (2) 是 p 阶相容的, 则它的局部截断误差满足

$$y_k - y(x_k) = O(h^{p+1}), \quad z_k - z(x_k) = O(h^p). \quad (23)$$

证明 令定理 1 中 $n = 0$, $\hat{y}_j = y(x_j)$, $\hat{z}_j = z(x_j)$, $j = 0, 1, \dots, k$, 这些值满足定理 1 的条件, 且 $\delta = O(h^p)$, $\theta = 0$, 由插值公式 (3) 知 $\|\hat{y}_0^h - y_0^h\| = O(h^{p+1})$, 应用定理 1 即得结论.

定理 2 设 2-DDAEs (1) 满足 $g_y f_z$ 可逆, $k \leq 6$, 方法 (2) 起始值满足

$$y_j - y(x_j) = O(h^{p+1}), \quad j = 0, 1, \dots, k-1, \quad (24)$$

则带 p 阶插值公式 (3) 的 k 步 BDF 方法 (2) 是 $p = k$ 阶收敛的, 即

$$y_n - y(x_n) = O(h^p), \quad z_n - z(x_n) = O(h^p), \quad x_n = nh, \quad n \geq k. \quad (25)$$

证明 首先研究 y 分量的局部误差积累. 数值解序列记为 $\{y_n^0, z_n^0\}$, 考虑多步解序列 $\{y_n^l, z_n^l\}$, $l = 1, 2, \dots$, 起始值为 $y_j^l = y(x_j)$, $z_j^l = z(x_j)$, $j = l-1, \dots, l+k-2$. 首要任务是估计 $y_n^l - y_n^{l+1}$, 为了方便起见, 我们去掉上标, 两个相邻多步解分别记为 $\{\hat{y}_n, \hat{z}_n\}$ 和 $\{\tilde{y}_n, \tilde{z}_n\}$. 为了能利用定理 1 的结论, 固定 3 个充分大的常数 $\hat{C}_0, \hat{C}_1, \hat{C}_2$ (文后注释说明了这 3 个常数的合理性), 使

$$\begin{aligned} \|\hat{y}_{n+j} - y(x_{n+j})\| &\leq \hat{C}_0 h, \quad \|\hat{z}_{n+j} - z(x_{n+j})\| \leq \hat{C}_1 h, \\ \|\hat{y}_{n+j} - \tilde{y}_{n+j}\| &\leq \hat{C}_2 h^2, \quad j = 0, 1, \dots, k-1. \end{aligned} \quad (26)$$

引入记号 $\Delta z_{n+k} = \tilde{z}_{n+k} - \hat{z}_{n+k}$, $\Delta y_{n+j} = \tilde{y}_{n+j} - \hat{y}_{n+j}$, $j = 0, 1, \dots, k$, $\Delta y_n^h = \tilde{y}_n^h - \hat{y}_n^h$, $\Delta Y_n = (\Delta y_{n+k-1}^T, \dots, \Delta y_n^T, (\Delta y_n^h)^T)^T$, 在定理 1 中取 $\delta = 0$, $\theta = 0$, 应用定理 1 可得

$$\|\Delta y_{n+k}\| \leq C(\|\Delta Y_n\|), \quad \|\Delta z_{n+k}\| \leq \frac{C}{h} \left(\sum_{j=0}^{k-1} \|g_y(\hat{y}_{n+k}) \Delta y_{n+j}\| + h \|\Delta Y_n\| \right). \quad (27)$$

只要 h 充分小时, 常数 C 不依赖于 $\hat{C}_0, \hat{C}_1, \hat{C}_2$ 的选取. 由插值公式 (3) 及假设 (26), 有

$$\|\Delta y_{n+k}\| = O(h^2), \quad \|\Delta z_{n+k}\| = O(h). \quad (28)$$

线性化多步公式得

$$\begin{aligned} \sum_{i=0}^k \alpha_i \Delta y_{n+i} &= h[f(\tilde{y}_{n+k}, \tilde{y}_n^h, \tilde{z}_{n+k}) - f(\hat{y}_{n+k}, \hat{y}_n^h, \hat{z}_{n+k})] \\ &= h f_z(\hat{y}_{n+j}, \hat{y}_n^h, \hat{z}_{n+j}) \Delta z_{n+j} + O(h \|\Delta Y_n\|), \end{aligned} \quad (29)$$

$$0 = g_y(\hat{y}_{n+k}) \Delta y_{n+k} + O(h \|\Delta Y_n\|). \quad (30)$$

记 $Q_{n+j} = (f_z(g_y f_z)^{-1} g_y)(\hat{y}_{n+j}, \hat{y}_n^h, \hat{z}_{n+j})$, $P_{n+j} = I - Q_{n+j}$, $j = 0, 1, \dots, k$, 易证

$$Q_{n+j}^2 = Q_{n+j}, \quad P_{n+j}^2 = P_{n+j}, \quad Q_{n+j} P_{n+j} = P_{n+j} Q_{n+j} = 0, \quad Q_{n+j+1} = Q_{n+j} + O(h), \quad (31)$$

用 P_{n+k} 左乘 (29), $f_z(g_y f_z)^{-1}$ 左乘 (30), 可得

$$\sum_{i=0}^k \alpha_i P_{n+i} \Delta y_{n+i} = O(h \|\Delta Y_n\|), \quad Q_{n+k} \Delta y_{n+k} = O(h \|\Delta Y_n\|). \quad (32)$$

由插值公式 (3), 延迟函数 $0 < \tau'(x) < 1$ 可导, 假设 (26) 及 (27), 有

$$\frac{1}{2} \Delta y_{n+1}^h = \frac{1}{2} \Delta y_n^h + O(h \|\Delta Y_n\|), \quad (33)$$

记

$$U_n = \left((P_{n+k-1} \Delta y_{n+k-1})^T, \dots, (P_n \Delta y_n)^T, \frac{1}{2} (\Delta y_n^h)^T \right)^T, \\ V_n = \left((Q_{n+k-1} \Delta y_{n+k-1})^T, \dots, (Q_n \Delta y_n)^T, \frac{1}{2} (\Delta y_n^h)^T \right)^T,$$

有 $\Delta Y_n = U_n + V_n$, (32) 式可分别化为

$$U_{n+1} = (A \otimes I) U_n + O(h \|U_n\| + h \|V_n\|), \quad (34)$$

$$V_{n+1} = (N \otimes I) V_n + O(h \|U_n\| + h \|V_n\|), \quad (35)$$

这里

$$A = \begin{pmatrix} -\alpha'_{k-1} & \cdots & -\alpha'_1 & -\alpha'_0 & 0 \\ 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 & 0 \\ 0 & \cdots & 0 & 0 & 1 \end{pmatrix}, \quad N = \begin{pmatrix} 0 & \cdots & 0 & 0 & 0 \\ 1 & \cdots & 0 & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & \cdots & 1 & 0 & 0 \\ 0 & \cdots & 0 & 0 & 1 \end{pmatrix},$$

其中 $\alpha'_j = \alpha_j / \alpha_k$. 由于 $\rho(\xi) = \sum_{j=0}^k \alpha_k \xi^k$ 满足根条件, 类似于文献 [18], 可选取一范数 $\|\cdot\|$, 使得 $\|A \otimes I\| \leq 1$, 及另一可能不同的范数 $\|\cdot\|$, 使得 $\|N \otimes I\| \leq 1$, 因此有

$$\begin{pmatrix} \|U_{n+1}\| \\ \|V_{n+1}\| \end{pmatrix} \leq \begin{pmatrix} 1 + O(h) & O(h) \\ O(h) & 1 + O(h) \end{pmatrix} \begin{pmatrix} \|U_n\| \\ \|V_n\| \end{pmatrix}, \quad (36)$$

化 (36) 中系数矩阵成对角阵, 易得

$$\begin{pmatrix} \|U_n\| \\ \|V_n\| \end{pmatrix} \leq T^{-1} \begin{pmatrix} \lambda_1^n & 0 \\ 0 & \lambda_2^n \end{pmatrix} T \begin{pmatrix} \|U_0\| \\ \|V_0\| \end{pmatrix}, \quad (37)$$

这里 $\lambda_1 = 1 + O(h)$, $\lambda_2 = 1 + O(h)$, 变换矩阵 T 由相应特征向量构成, 满足

$$T = \begin{pmatrix} 1 & O(h) \\ O(h) & 1 \end{pmatrix},$$

化简得

$$\|U_n\| \leq \check{C}(\|U_0\| + h\|V_0\|), \quad (38)$$

$$\|V_n\| \leq \check{C}(h\|U_0\| + \|V_0\|), \quad (39)$$

$$\|Y_n\| \leq C_6(\|U_n\| + \|V_n\|) \leq C_7(\|U_0\| + \|V_0\|). \quad (40)$$

向量 U_0, V_0 由 y 变量的起始值的误差及局部截断误差组成, 满足

$$\|U_0\| \leq H_0 h^{p+1}, \quad \|V_0\| \leq H_1 h^{p+1}, \quad \|\Delta y_0\| \leq H_2 h^{p+1}, \quad (41)$$

由 (23), (24), (38)-(40) 及 (41), 可知

$$\|\Delta y_n\| \leq C_8 h^{p+1}, \quad \|g_y(\hat{y}_{n+k})\Delta y_{n+j}\| \leq C_9 h^{p+1}, \quad (42)$$

$$\|y_n - y(x_n)\| \leq \sum_{l=0}^{n-k+1} \|y_n^l - y_n^{l+1}\| \leq C_{10} h^p. \quad (43)$$

由于 z_n 仅依赖于 $y_{n-k}^h, y_{n-k}, \dots, y_{n-1}$, 在定理 1 中取 $\hat{y}_i = y(x_i)$, $\hat{z}_i = z(x_i)$, $\delta = O(h^p)$, $\theta = 0$, 应用定理 1 的结论可得

$$\|z_n - z(x_n)\| \leq \frac{C}{h} \sum_{j=0}^k \|g_y(y(x_n))(y_{n-j} - y(x_{n-j}))\| + O(h^p). \quad (44)$$

利用 (42) 和 (43), 可得

$$\begin{aligned} \|g_y(y(x_n))(y_{n-j} - y(x_{n-j}))\| &= \sum_{l=0}^{n-k+1} \|g_y(y_n^l + O(h^p))(y_{n-j}^l - y_{n-j}^{l+1})\| \\ &\leq \sum_{l=0}^{n-k+1} (\|g_y(y_n^l)(y_{n-j}^l - y_{n-j}^{l+1})\| + O(h^{2p+1})) = O(h^{p+1}), \end{aligned}$$

因此

$$\|z_n - z(x_n)\| \leq C_{11} h^p. \quad (45)$$

注 1 一般来说, 常数 C_{10}, C_{11} 依赖于假设 (26) 中的 \hat{C}_0, \hat{C}_1 , 我们严格限制步长 h , 使得 $C_{10} h^{p-1} \leq \hat{C}_0$, $C_{11} h^{p-1} \leq \hat{C}_1$, 这样, 数值解就不会与假设 (26) 相冲突.

3 数值试验

考虑如下 2-指标变延迟微分代数初值问题

$$\begin{cases} y_1'(x) = -2\sqrt{y_1 z} + y_2 \frac{y_1(\frac{x}{2})}{y_2(\frac{x}{2})}, & 0 \leq x \leq 2, \\ y_2'(x) = -\frac{z^2}{y_2} - y_1 y_2(\frac{x}{2}) - \sqrt{z^3}, & 0 \leq x \leq 2, \\ 0 = y_1 y_2 + y_2^2, & 0 \leq x \leq 2, \\ y_1(0) = 1, \quad y_2(0) = -1, \quad z(0) = 1, \end{cases} \quad (46)$$

问题 (46) 有唯一真解 $y_1(x) = e^{-x}$, $y_2(x) = -e^{-x}$, $z(x) = e^{-x}$. 取步长 h , y_1, y_2, z 在 $x = 2$ 处的整体误差分别记为 $yerr1(h)$, $yerr2(h)$, $zerr(h)$. 通过计算可以得到相应的误差阶, 分别记为 $py1(h)$, $py2(h)$, $pz(h)$, 其中

$$py1(h) = \ln \frac{yerr1(h)}{yerr1(0.5h)} / \ln 2, \quad py2(h) = \ln \frac{yerr2(h)}{yerr2(0.5h)} / \ln 2, \quad pz(h) = \ln \frac{zerr(h)}{zerr(0.5h)} / \ln 2.$$

分别用带线性插值公式的二步二阶 BDF 方法 (BDF2) 和带二次插值公式的三步三阶 BDF 方法 (BDF3) 求解初值问题 (46). 数值结果如表 1 和表 2 所示, 验证了理论结果的正确性.

表 1: BDF2 的数值结果

h	$yerr1$	$yerr2$	$zerr$	$py1$	$py2$	pz	CPU
0.1	$0.3590E-2$	$0.4899E-2$	$0.7199E-2$	1.9974	2.0283	2.0865	0.1590
0.05	$0.8991E-3$	$0.1201E-2$	$0.1695E-2$	2.0031	1.9750	1.9924	0.3167
0.025	$0.2243E-3$	$0.3055E-3$	$0.4260E-3$	2.0019	1.9976	2.0122	0.6085
0.0125	$0.0560E-3$	$0.0765E-3$	$0.1056E-3$				1.1973

表 2: BDF3 的数值结果

h	$yerr1$	$yerr2$	$zerr$	$py1$	$py2$	pz	CPU
0.1	$0.5100E-3$	$0.7001E-3$	$0.1100E-2$	2.9521	2.9436	3.0179	0.2179
0.05	$0.6590E-4$	$0.9100E-4$	$0.1358E-3$	2.9684	3.0019	3.0541	0.4064
0.025	$0.8420E-5$	$0.1136E-4$	$0.1635E-4$	3.0076	2.9501	3.0589	0.7932
0.0125	$0.1047E-5$	$0.1470E-5$	$0.1962E-5$				1.4571

参考文献:

[1] 余越昕, 文立平, 李寿佛. 延迟积分微分方程单支方法的稳定性分析[J]. 工程数学学报, 2008, 25(3): 469-474
Yu Y X, Wen L P, Li S F. Stability analysis of one-leg methods of delayed integro-differential equations[J]. Chinese Journal of Engineering Mathematics, 2008, 25(3): 469-474

[2] 倪涛洋, 黎培兴, 李彩霞. 中立型线性微分方程的稳定性[J]. 工程数学学报, 2005, 22(1): 144-146
Ni T Y, Li P X, Li C X. The stability of neutral linear differential equation[J]. Chinese Journal of Engineering Mathematics, 2005, 22(1): 144-146

[3] Petzold L R. Differential-algebraic equations are not ODEs[J]. SIAM Journal on Scientific and Statistical Computing, 1982, 3(3): 367-384

[4] Zhu W L, Petzold L R. Asymptotic stability of linear differential-algebraic equations and numerical methods[J]. Applied Numerical Mathematics, 1997, 24(2): 247-264

[5] Campbell S L, Linh V H. Stability criteria for differential-algebraic equations with multiple delays and their numerical solutions[J]. Applied Mathematics and Computation, 2009, 208(2): 397-415

[6] Shampine L F, Gahinet P. Delay differential-algebraic equations in control theory[J]. Applied Numerical Mathematics, 2006, 56(3): 574-588

[7] Baker C T H, Paul C A H, Tian H. Differential algebraic equations with after-effect[J]. Journal of Computational and Applied Mathematics, 2002, 140(1): 63-80

[8] Xu Y, Zhao J J, Dong S Y, et al. Stability of the Rosenbrock methods for the neutral delay differential-algebraic equations[J]. Applied Mathematics and Computation, 2005, 168(2): 1128-1144

- [9] Xu Y, Zhao J J, Sui Z N. Stability analysis of θ -methods for neutral multi-delay-integro-differential system[J]. Discrete Dynamics in Nature and Society, 2008, 79(1): 571-583
- [10] 张诚坚, 程玮. 延迟微分代数系统的隐式中点法稳定性判据[J]. 华中理工大学学报, 2000, 28(2): 112-113
Zhang C J, Cheng W. The stability criteria of implicit midpoint rule for delay differential-algebraic systems[J]. Journal of Huazhong University of Science and Technology, 2000, 28(2): 112-113
- [11] 张诚坚, 廖晓昕. (ρ, σ) -方法关于刚性延迟微分代数系统的非线性稳定性[J]. 控制理论与应用, 2001, 18(3): 827-832
Zhang C J, Liao X Q. Nonlinear stability of (ρ, σ) -methods for stiff delay-differential-algebraic systems[J]. Control Theory & Applications, 2001, 18(3): 827-832
- [12] 李宏智. 延迟微分代数系统的数值稳定性及其块方法[D]. 武汉: 华中科技大学, 2004
Li H Z. The numerical stability and block methods for delay-differential-algebraic systems[D]. Wuhan: Huazhong University of Science and Technology, 2004
- [13] 甘四清. 奇异摄动问题与时滞奇异摄动问题数值分析[D]. 北京: 中国科学院数学与系统科学研究院数学研究所, 2001
Gan S Q. The numerical analysis for singular perturbation and delay singular perturbation problems[D]. Beijing: Institute of Mathematics and System Science, Chinese Academy of Sciences, 2001
- [14] Ascher U M, Petzold L R. The numerical solution of delay-differential-algebraic equations of retarded and neutral type[J]. SIAM Journal on Numerical Analysis, 1995, 32(5): 1635-1657
- [15] Hauber R. Numerical treatment of retarded differential-algebraic equations by collocation methods[J]. Advances in Computational Mathematics, 1997, 7(4): 247-264
- [16] 肖飞雁, 张诚坚. 一类变时滞微分代数方程单支方法的收敛性[J]. 数值计算与计算机应用, 2008, 29(3): 217-225
Xiao F Y, Zhang C J. Convergence of one-leg methods for a class of variable retarded differential algebraic equations[J]. Journal on Numerical Methods and Computer Applications, 2008, 29(3): 217-225
- [17] Xiao F Y, Zhang C J. Convergence analysis of Runge-Kutta methods for a class of retarded differential algebraic systems[J]. Acta Mathematica Scientia, 2010, 30(1): 65-74
- [18] Hairer E, Wanner G. Solving Ordinary Differential Equations II: Stiff and Differential-Algebraic Problems[M]. Berlin: Springer-Verlag, 1996

Convergence of Backward Differentiation Formulas for Index-2 Differential-algebraic Equations with Variable Delay

LIU Hong-liang, XIAO Ai-guo

(School of Mathematics and Computational Science, Hunan Key Laboratory for Computation and Simulation in Science and Engineering, Xiangtan University, Xiangtan 411105)

Abstract: Delay differential-algebraic equations often arise in automatic control, power and circuit analysis, multi-body dynamics, etc. The current researches on numerical analysis for delay-differential-algebraic equations are mainly focused on linear problems and 1-index problems. It is difficult to do numerical analysis for high-index nonlinear delay-differential-algebraic equations, and there are only a few results for this kind of problems, furthermore, most of them are about constant-delay problems. The backward differentiation formulas are applied in this paper to index-2 nonlinear differential-algebraic equations with variable delay. The corresponding convergence results are obtained and confirmed by some numerical examples.

Keywords: index-2 differential-algebraic equations; backward differentiation formulas; convergence; variable delay

Received: 05 Aug 2009. **Accepted:** 14 Dec 2010.

Foundation item: The National Natural Science Foundation of China (10971175); the Specialized Research Fund for the Doctoral Program of Higher Education of China (20094301110001); the Natural Science Foundation of Hunan Province (09JJ3002); the Doctoral Research Fund of Xiangtan University (11QDZ01).